

Fish Belt Transect Form
魚類帶斷面表

Site Name 調查區名稱:					
Average Depth 平均深度:		Team Leader 調查隊長:			
Date 日期:		Time 時間:			
Indo-Pacific (Hong Kong) Belt Transect : Fish 魚類					
Data recorded by 資料記錄者:					
		0-20m	25-45m	50-70m	75-95m
Butterfly fish 蝴蝶魚					
Sweetlips (Haemulidae) 細鱗					
Grouper 石斑魚:					
Barramundi cod 老鼠斑 (<i>Cromileptes altivelis</i>)	>30 cm				
	<30 cm				
Coral Trout 星斑 (<i>Plectropomus</i> sp.)	>30 cm				
	<30 cm				
Other Grouper 其他石斑	>30 cm				
	<30 cm				
Snappers 笛鯛類 (Lutjanidae)	>30 cm				
	<30 cm				
Wrasse 眉魚:					
Humphead wrasse (<i>Cheilinus undulatus</i>) □□					
Others Wrasse (Labridae) 雜眉					
Parrotfish 鸚鵡魚:					
Bumphead parrot 駝背大鸚鵡魚					
Blue-barred parrot (<i>Scarus ghobban</i>) 黃衣					
Other Parrotfish 其他鸚鵡魚					
Seahorse 海馬					
Moray Eel 油 鯧					
Other Notes 其他記錄		0-20m	25-45m	50-70m	75-95m
Bleaching (% of coral population and % per colony): 白化 (佔珊瑚群體的%□□□□□□□%):					
Suspected disease (type & %) 病態 (類型及%)					
Rare animals sighted (type & no.) 罕見生物 (類型及□□)					